



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

A LIST OF BIRDS TAKEN ON MARAGUANA,  
WATLING'S ISLAND, AND INAGUA, BAHAMA,  
DURING JULY, AUGUST, SEPTEMBER,  
AND OCTOBER, 1891.

BY CHARLES B. CORY.

MARAGUANA (August 5 to September 14).

<i>Larus atricilla</i> Linn.	<i>Ceryle alcyon</i> (Linn.).
<i>Gelochelidon nilotica</i> (Hasselq.).	<i>Tyrannus dominicensis</i> (Gmel.).
<i>Ardea tricolor ruficollis</i> Gosse.	<i>Loxigilla violacea</i> (Linn.).
<i>Ardea virescens</i> (Linn.).	<i>Spindalis zena</i> (Linn.).
<i>Nycticorax violaceus</i> (Linn.).	<i>Euethia bicolor</i> (Linn.).
<i>Himantopus mexicanus</i> (Müll.).	<i>Chelidon erythrogaster</i> (Bodd.).
<i>Tringa maculata</i> Vieill.	<i>Vireo crassirostris</i> (Bryant).
<i>Tringa minutilla</i> Vieill.	<i>Cœreba bahamensis</i> (Reich.).
<i>Totanus flavipes</i> (Gmel.).	<i>Dendroica petechia gundlachi</i>
<i>Symphemia semipalmata</i> (Gmel.).	Baird.
<i>Bartramia longicauda</i> (Bechst.).	<i>Dendroica dominica</i> (Linn.).
<i>Ægialitis vocifera</i> (Linn.).	<i>Dendroica discolor</i> (Vieill.).
<i>Ægialitis wilsonia</i> (Ord).	<i>Mimus polyglottus orpheus</i>
<i>Arenaria interpres</i> (Linn.).	(Linn.).
<i>Hæmatopus palliatus</i> Temm.	<i>Mimus gundlachi</i> Caban.
<i>Columba leucocephala</i> (Linn.).	<i>Polioptila cærulea</i> (Linn.).
<i>Columbigallina passerina</i> (Linn.).	<i>Margarops fuscatus</i> (Vieill.).

WATLING'S ISLAND (September 28 to November 9).

<i>Larus atricilla</i> (Linn.).
<i>Phalacrocorax dilophus floridanus</i> (Aud.). — Eight specimens.
<i>Ardea tricolor ruficollis</i> (Gosse).
<i>Ægialitis vocifera</i> (Linn.).
<i>Coccyzus minor maynardi</i> Ridgw.
<i>Coccyzus americanus</i> (Linn.).
<i>Tyrannus dominicensis</i> (Gmel.).
<i>Dolichonyx oryzivorus</i> (Linn.). — Oct. 12, female.
<i>Habia ludoviciana</i> (Linn.). — Oct. 20, female.
<i>Euethia bicolor</i> (Linn.).
<i>Vireo olivaceus</i> (Linn.). — Oct. 5, male.
<i>Cœreba bahamensis</i> (Reich.).
<i>Mniotilta varia</i> (Linn.).
<i>Compsothlypis americana</i> (Linn.). — Five specimens.

- Dendroica tigrina* (Gmel.). — Seven specimens.  
*Dendroica petechia gundlachi* Baird. — Common, breeds.  
*Dendroica maculosa* (Gmel.). — Six specimens, Oct. 6-21.  
*Dendroica striata* (Forst.). — Thirty-six specimens.  
*Dendroica blackburniæ* (Gmel.). — Two specimens, Oct. 12.  
*Dendroica dominica* (Linn.). — Six specimens.  
*Dendroica virens* (Gmel.). — Oct. 12, male.  
*Dendroica palmarum* (Gmel.). — Oct. 20, female.  
*Dendroica discolor* (Vieill.). — Twenty-five specimens.  
*Seiurus aurocapillus* (Linn.). — Thirteen specimens.  
*Mimus gundlachi* Caban.  
*Margarops fuscatus* (Vieill.).

## INAGUA.

- Gallinula galeata* (Licht.). — July 17.  
*Tringa minutilla* Vieill. — July 28.  
*Totanus flavipes* (Gmel.). — July 28.  
*Myiarchus sagræ* Gundl.  
*Loxigilla violacea* (Linn.).  
*Vireo olivaceus* (Linn.). — Sept. 17, female.  
*Vireo crassirostris* (Bryant).  
*Cœreba bahamensis* (Reich.).  
*Helmitherus vermivorus* (Gmel.). — Sept. 22, female.  
*Dendroica petechia gundlachi* Baird.  
*Mimus gundlachi* Caban.

---

## NOTES ON THE RANGE AND HABITS OF THE CAROLINA PARRAKEET.

BY AMOS W. BUTLER.

THE CAROLINA PARRAKEET (*Conurus carolinensis*), whose range is now confined to quite restricted areas in some of our southern States, was formerly known as a characteristic bird of Indiana. At the time of its greatest range in that State, within historic times, it was known from New York, Pennsylvania and Maryland to Kansas, Nebraska, and possibly Colorado. It is my desire to present some evidence tending to show its distribution in Indiana and neighboring States together with some notes upon its habits.